
## Directions for filling out EMERGENCY CARE PLAN FOR ANAPHYLACTIC ALLERGIES form.


- 1. Use Adobe Acrobat Reader to view form.
  - Click link below if download of Adobe Acrobat Reader is necessary.



- 2. Please **DOWNLOAD** and **OPEN** file locally before completing by clicking the download icon in browser being used.
  - Firefox browser



• Google Chrome browser



• Internet Explorer browser



3. Please submit the completed form to the School or Nurse's Office.

Madeline Brick, R.N.

Phone: 671-477-6341 ext. 280

Fax: 671-477-7136

madeline.brick@hbcguam.net



## **EMERGENCY CARE PLAN FOR ANAPHYLACTIC ALLERGIES**

Last Name	First Name	Grade Teacher
Allergens:		Birthdate
Has epi-pen ever been used in the past? Yes No No If yes, when?		
Asthmatic? Yes No Inhaler at school? Yes No		
Father's Last Name		
		Work Phone
Mother's Last Name		
Home Phone	Cell Phone	Work Phone
Emergency Contact Name	Relationship	Contact No.
Emergency Contact Name	Relationship	Contact No.
Symptoms of an allergic reaction may include any/all of these: (Check appropriate boxes.)		
► Mouth Itching & swelling of lips	Itching & swelling of tongue or mouth	Mouth "feels hot"
► Throat Tightness in throat	Itching Hoarseness C	Cough Difficulty swallowing
► <b>Skin</b> Swelling of face & extremities	Itchy rash Hives	
► Stomach Nausea/Vomiting	Abdominal cramps Diarrh	nea 🔲
► Lung Shortness of breath	Repetitive cough Wheez	zing
► <b>Heart</b> "Thready pulse"	Dizzy/"Passing out" Pale	
Staff members notified of allergy: Homeroom Teacher		
TREATMENT PLAN FOR SCHOOL STAFF TO FOLLOW (To be filled out by parents or doctor.)		
► Call school nurse. ► Call Parent/guardian as soon as possible. ► If ingestion or suspected ingestion of allergen occurs and symptoms are present and epinephrine is ordered, give epinephrine immediately and call 911.		
Treatment should be initiated with symptoms		- · · · · · · · · · · · · · · · · · · ·
Benadryl ordered Yes No	Dosage of Bo	enadryl per provider's orders
Epinephrine ordered Yes No	Special instructions	
Location of Epinephrine Nurse's office	Main Classroom With studen	other
Expiration Date of Epinephrine		
Preferred hospital if transported		
Doctor's Name Contact No		
Epinephrine provides a 20 minute response window. After epinephrine, a student may feel dizzy or have an increased heart rate. This is a normal response. Students receiving epinephrine should be transported to the hospital by ambulance. A staff member should accompany the student to the emergency room if the parent, guardian, or emergency contact is not present and adequate supervision for others is present.		
HEALTHCARE PROVIDER:		PHONE:

PARENT'S/GUARDIAN'S SIGNATURE: \_\_\_\_\_ DATE: \_\_\_\_\_